

# **Differential Prognostic Impact of Treatment Strategies for LM versus Non-LM Bifurcation Lesion: Lessons from the COBIS Registries**

**Young Bin Song, MD**  
**On the Behalf of COBIS III Investigators**



**Heart Vascular Stroke Institute,  
Samsung Medical Center, Seoul, Republic of Korea**

# Coronary Bifurcation Stent (COBIS) Registries

- Multi-center retrospective real-world registry of drug-eluting stenting in coronary bifurcation lesions in Korea

	<b>COBIS I</b>	<b>COBIS II</b>	<b>COBIS III</b>
<b>Enrolled Patients</b>	1691	2897	2648
<b>Term</b>	2004.1~2006.6	2003.1 ~ 2009.12	2010.1 ~ 2014.12
<b>Main vessel Diameter (mm)</b>	≥ 2.5	≥ 2.5	≥ 2.5
<b>Side branch Diameter (mm)</b>	≥ 2.0	≥ 2.3	≥ 2.3
<b>LM Bifurcation</b>	X	0	0
<b>DES generation</b>	1 <sup>st</sup>	1 <sup>st</sup> + 2 <sup>nd</sup> (14%)	2 <sup>nd</sup> only
<b>Publications</b>	12	19	-


# Stent Technique (1-Stent vs. 2-Stent)




# Stent Technique (1-Stent VS. 2-Stent)

## LM vs. Non LM Bifurcation Lesions from COBIS II Registry - Target Lesion Failure -

Non-LM




LM



	Non-LM Bifurcation (n = 2,044)		LM Bifurcation (n = 853)		Interaction p Value
	HR (95% CI)	p Value	HR (95% CI)	p Value	
Target lesion failure	1.39 (0.99–1.94)	0.06	2.38 (1.60–3.55)	<0.01	<0.01


# Stent Technique (1-Stent vs. 2-Stent)

## LM vs. Non LM Bifurcation Lesions from COBIS II Registry - Cardiac Death or MI -



The **1-stent strategy**, if possible, should initially be considered the preferred approach for the treatment of coronary bifurcation lesions, **especially LM bifurcation lesions**.

# Study Flow



**Primary Endpoint**

**Target Lesion Failure, a composite of cardiac death, myocardial infarction, and target lesion revascularization**

# Baseline Clinical Characteristics


Variables	Left Main Bifurcation (N=935)			Non-Left Main Bifurcation (N=1713)		
	1-Stent (N=682)	2-Stent (N=253)	P value	1-Stent (N=1512)	2-Stent (N=201)	P value
<b>Demographics</b>						
Age (years)	65.0 ± 10.4	66.8 ± 11.3	0.022	62.8 ± 11.2	62.8 ± 10.4	0.943
Male sex	524 (76.8%)	187 (73.9%)	0.399	1158 (76.6%)	144 (71.6%)	0.146
<b>Cardiovascular risk factors</b>						
Hypertension	419 (61.4%)	138 (54.5%)	0.067	831 (55.0%)	116 (57.7%)	0.508
Diabetes mellitus	262 (38.4%)	94 (37.2%)	0.782	<b>467 (30.9%)</b>	<b>82 (40.8%)</b>	<b>0.006</b>
Chronic kidney disease	38 (5.6%)	9 (3.6%)	0.278	45 (3.0%)	11 (5.5%)	0.097
Hyperlipidemia	<b>280 (41.1%)</b>	<b>81 (32.0%)</b>	<b>0.014</b>	<b>555 (36.7%)</b>	<b>93 (46.3%)</b>	<b>0.011</b>
Current smoking	172 (25.2%)	54 (21.3%)	0.253	507 (33.5%)	65 (32.3%)	0.797
Previous PCI	114 (16.7%)	45 (17.8%)	0.772	143 (9.5%)	21 (10.4%)	0.748
Previous myocardial infarction	35 (5.1%)	13 (5.1%)	>0.999	57 (3.8%)	8 (4.0%)	>0.999
Previous CVA	57 (8.4%)	19 (7.5%)	0.774	93 (6.2%)	8 (4.0%)	0.285
<b>Initial presentation</b>						
Clinical presentation			0.626			0.179
Stable ischemic heart disease	277 (40.6%)	100 (39.5%)		573 (37.9%)	79 (39.3%)	
Unstable angina or NSTEMI	351 (51.5%)	137 (54.2%)		733 (48.5%)	104 (51.7%)	
STEMI	54 (7.9%)	16 (6.3%)		206 (13.6%)	18 (9.0%)	
LVEF (%)	58.5 ± 10.2	57.6 ± 10.3	0.241	58.7 ± 9.7	59.5 ± 8.8	0.347
<b>Medications at discharge</b>						
Aspirin	667 (98.1%)	244 (96.8%)	0.365	1493 (98.7%)	198 (98.5%)	>0.999
P2Y12 inhibitors	672 (98.8%)	244 (96.8%)	0.072	1494 (98.8%)	198 (98.5%)	0.980
Beta-blocker	420 (61.6%)	173 (68.4%)	0.066	900 (59.5%)	127 (63.2%)	0.358
ACE inhibitor or ARB	407 (59.7%)	149 (58.9%)	0.887	940 (62.2%)	127 (63.2%)	0.840
Statin	600 (88.0%)	225 (88.9%)	0.773	1363 (90.1%)	181 (90.0%)	>0.999

# Baseline Lesion Characteristics

Variables	Left Main Bifurcation (N=935)			Non-Left Main Bifurcation (N=1713)		
	1-Stent (N=682)	2-Stent (N=253)	P value	1-Stent (N=1512)	2-Stent (N=201)	P value
<b>Lesion characteristics</b>						
Multi-vessel disease	532 (78.0%)	242 (95.7%)	<0.001	770 (50.9%)	103 (51.2%)	0.992
<b>Bifurcation location</b>						
LAD/diagonal				<b>1037 (68.6%)</b>	<b>171 (85.1%)</b>	
LCX/OM				326 (21.6%)	24 (11.9%)	
RCA (PL/PDA)				149 (9.9%)	6 (3.0%)	
Medina classification			<0.001			<0.001
1.1.1	93 (13.6%)	128 (50.6%)		514 (34.0%)	104 (51.7%)	
1.0.1	19 (2.8%)	19 (7.5%)		116 (7.7%)	14 (7.0%)	
0.1.1	29 (4.3%)	44 (17.4%)		126 (8.3%)	49 (24.4%)	
1.0.0	77 (11.3%)	7 (2.8%)		212 (14.0%)	0 (0%)	
1.1.0	164 (24.0%)	15 (5.9%)		241 (15.9%)	7 (3.5%)	
0.1.0	275 (40.3%)	9 (3.6%)		283 (18.7%)	8 (4.0%)	
0.0.1	25 (3.7%)	31 (12.3%)		20 (1.3%)	19 (9.5%)	
<b>True bifurcation</b>	<b>141 (20.7%)</b>	<b>191 (75.5%)</b>	<b>&lt;0.001</b>	<b>756 (50.0%)</b>	<b>167 (83.1%)</b>	<b>&lt;0.001</b>
<b>Procedural characteristics</b>						
<b>No. of used stent</b>	<b>1.7 ± 0.9</b>	<b>2.6 ± 1.0</b>	<b>&lt;0.001</b>	<b>1.6 ± 0.9</b>	<b>2.3 ± 1.1</b>	<b>&lt;0.001</b>
Stent type			0.161			0.011
Everolimus-eluting stents	367 (53.8%)	131 (51.8%)		668 (44.2%)	110 (54.7%)	
Zotarolimus-eluting stents	164 (24.0%)	69 (27.3%)		446 (29.5%)	57 (28.4%)	
Biolimus-eluting stents	132 (19.4%)	40 (15.8%)		317 (21.0%)	25 (12.4%)	
Mixed or other stents	19 (2.8%)	13 (5.1%)		81 (5.4%)	9 (4.5%)	
<b>Trans-radial intervention</b>	<b>366 (53.7%)</b>	<b>106 (41.9%)</b>	<b>0.002</b>	<b>963 (63.7%)</b>	<b>72 (35.8%)</b>	<b>&lt;0.001</b>
<b>IVUS guidance</b>	<b>427 (62.6%)</b>	<b>172 (68.0%)</b>	<b>0.148</b>	<b>389 (25.7%)</b>	<b>75 (37.3%)</b>	<b>0.001</b>
<b>Final kissing ballooning</b>	<b>163 (23.9%)</b>	<b>233 (92.1%)</b>	<b>&lt;0.001</b>	<b>228 (15.1%)</b>	<b>165 (82.1%)</b>	<b>&lt;0.001</b>
<b>POT</b>	<b>237 (34.8%)</b>	<b>56 (22.1%)</b>	<b>&lt;0.001</b>	<b>394 (26.1%)</b>	<b>52 (25.9%)</b>	<b>&gt;0.999</b>
<b>Re-POT</b>	<b>25 (3.7%)</b>	<b>48 (19.0%)</b>	<b>&lt;0.001</b>	<b>23 (1.5%)</b>	<b>27 (13.4%)</b>	<b>&lt;0.001</b>
<b>NC balloon use</b>	<b>162 (23.8%)</b>	<b>87 (34.4%)</b>	<b>0.001</b>	<b>228 (15.1%)</b>	<b>57 (28.4%)</b>	<b>&lt;0.001</b>


# Procedural Characteristics

## Stent Technique (Type of 2-Stent Strategy)




# Clinical Outcomes Between LM VS Non-LM


**A. Target Lesion Failure**



**B. Cardiac Death or Myocardial Infarction**



**C. Target Lesion Revascularization**




**Patients with LM bifurcation showed significantly higher risks of target lesion failure, cardiac death or myocardial infarction, and target lesion revascularization compared to those with non-LM bifurcation.**




# Clinical Outcomes Between 1-Stent vs 2-Stent in LM Bifurcation


**A. Target Lesion Failure**






**B. Cardiac Death or Myocardial Infarction**



**C. Target Lesion Revascularization**




## Comparison with COBIS II Registry






# Clinical Outcomes Between 1-Stent vs 2-Stent in Non-LM Bifurcation


### A. Target Lesion Failure






### B. Cardiac Death or Myocardial Infarction



### C. Target Lesion Revascularization



## Comparison with COBIS II Registry




# Association Between Treatment Strategy and Bifurcation Location

Outcomes	Groups	Multivariable Adjustment*				IPW Analysis†			
		HR	95% CI	P value	Interaction P	HR	95% CI	P value	Interaction P
TLF‡	All	1.368	0.888-2.107	0.156	<b>0.045</b>	1.531	1.007-2.328	0.046	<b>0.020</b>
	<b>LM</b>	<b>1.848</b>	<b>1.045-3.266</b>	<b>0.035</b>		<b>2.332</b>	<b>1.388-3.918</b>	<b>0.001</b>	
	Non-LM	0.925	0.428-2.001	0.843		0.730	0.319-1.670	0.456	
Cardiac Death or MI	All	0.687	0.361-1.310	0.255	0.885	0.672	0.328-1.380	0.279	0.933
	LM	0.991	0.413-2.379	0.984		0.656	0.306-1.407	0.279	
	Non-LM	0.588	0.203-1.705	0.329		0.700	0.216-2.265	0.551	
All-cause Death	All	1.033	0.565-1.888	0.917	0.271	0.962	0.494-1.872	0.908	0.405
	LM	1.645	0.639-4.231	0.302		0.711	0.303-1.669	0.433	
	Non-LM	0.928	0.404-2.134	0.860		1.193	0.488-2.917	0.699	
Cardiac Death	All	0.727	0.315-1.679	0.455	0.185	0.755	0.296-1.926	0.556	0.276
	LM	1.001	0.280-3.584	0.999		0.433	0.144-1.301	0.136	
	Non-LM	0.856	0.271-2.708	0.791		1.051	0.319-3.460	0.934	
MI	All	0.703	0.270-1.834	0.472	0.297	0.627	0.261-1.503	0.295	0.421
	LM	0.709	0.212-2.374	0.577		0.848	0.304-2.363	0.752	
	Non-LM	0.399	0.044-3.669	0.417		0.343	0.046-2.551	0.296	
TLR	All	2.333	1.307-4.165	0.004	<b>0.015</b>	2.305	1.395-3.800	0.001	<b>0.007</b>
	<b>LM</b>	<b>3.062</b>	<b>1.448-6.477</b>	<b>0.003</b>		<b>3.596</b>	<b>1.954-6.619</b>	<b>&lt;0.001</b>	
	Non-LM	1.214	0.404-3.642	0.730		0.655	0.226-1.902	0.437	


# Comparison Between COBIS II and COBIS III Registries

	COBIS II (2003-2009)	COBIS III (2010-2014)
<b>Enrolled Patients</b>	<b>2897</b>	<b>2648</b>
<b>Median Follow-up, Months</b>	<b>36 (25-52)</b>	<b>53 (37-68)</b>
<b>LM Bifurcation</b>	<b>853 (29.4%)</b>	<b>935 (35.3%)</b>
<b>Second-generation DES</b>	<b>422 (14.6%)</b>	<b>2648 (100%)</b>
<b>2-Stent Strategy</b>	<b>770 (26.6%)</b>	<b>454 (17.1%)</b>
<b>Cumulative Incidence of Target Lesion Failure at 5 years</b>	<b>286 (12.6%)</b>	<b>184 (8.4%)</b>

**A. Target Lesion Failure in LM Bifurcation**




**B. Target Lesion Failure in Non-LM Bifurcation**




# Conclusion and Summary

## A. Left Main Bifurcation



## B. Non-Left Main Bifurcation



1. Patients treated with PCI for an **LM bifurcation lesion** had **poorer outcomes** than those with a non-LM bifurcation lesion in the era of second-generation DES.
2. In the treatment of LM bifurcation, the **2-stent strategy** was associated with a **higher risk of TLF** than the 1-stent strategy, mainly driven by higher risk of TLR; however, **incidence of cardiac death or MI was not different between the strategies**, unlike previous results from the COBIS II registry.
3. In the treatment of a **non-LM bifurcation**, there were **no significant differences in TLF, cardiac death or MI, and TLR** between the 1-stent and 2-stent strategy groups.

# **Thank You For Your Attention !**

**Young Bin Song, MD, PhD**

**Heart Vascular Stroke Institute,  
Samsung Medical Center, Seoul, Republic of Korea**